

COVID-19 vaccine efficacy and effectiveness—the elephant (not) in the room



Approximately 96 COVID-19 vaccines are at various stages of clinical development.¹ At present, we have the interim results of four studies published in scientific journals (on the Pfizer–BioNTech BNT162b2 mRNA vaccine,² the Moderna–US National Institutes of Health [NIH] mRNA-1273 vaccine,³ the AstraZeneca–Oxford ChAdOx1 nCov-19 vaccine,⁴ and the Gamaleya GamCovidVac [Sputnik V] vaccine)⁵ and three studies through the US Food and Drug Administration (FDA) briefing documents (on the Pfizer–BioNTech,⁶ Moderna–NIH,⁷ and Johnson & Johnson [J&J] Ad26.COV2.S vaccines).⁸ Furthermore, excerpts of these results have been widely communicated and debated through press releases and media, sometimes in misleading ways.⁹ Although attention has focused on vaccine efficacy and comparing the reduction of the number of symptomatic cases, fully understanding the efficacy and effectiveness of vaccines is less straightforward than it might seem. Depending on how the effect size is expressed, a quite different picture might emerge (figure; appendix).

Vaccine efficacy is generally reported as a relative risk reduction (RRR). It uses the relative risk (RR)—ie, the ratio of attack rates with and without a vaccine—which is expressed as 1–RR. Ranking by reported efficacy gives relative risk reductions of 95% for the Pfizer–BioNTech, 94% for the Moderna–NIH, 90% for the Gamaleya, 67% for the J&J, and 67% for the AstraZeneca–Oxford vaccines. However, RRR should be seen against the background risk of being infected and becoming ill with COVID-19, which varies between populations and over time. Although the RRR considers only participants who could benefit from the vaccine, the absolute risk reduction (ARR), which is the difference between attack rates with and without a vaccine, considers the whole population. ARR tends to be ignored because they give a much less impressive effect size than RRRs: 1.3% for the AstraZeneca–Oxford, 1.2% for the Moderna–NIH, 1.2% for the J&J, 0.93% for the Gamaleya, and 0.84% for the Pfizer–BioNTech vaccines.

ARR is also used to derive an estimate of vaccine effectiveness, which is the number needed to vaccinate (NNV) to prevent one more case of COVID-19 as 1/ARR.

NNVs bring a different perspective: 76 for the Moderna–NIH, 78 for the AstraZeneca–Oxford, 80 for the Gamaleya, 84 for the J&J, and 117 for the Pfizer–BioNTech vaccines. The explanation lies in the combination of vaccine efficacy and different background risks of COVID-19 across studies: 0.9% for the Pfizer–BioNTech, 1% for the Gamaleya, 1.4% for the Moderna–NIH, 1.8% for the J&J, and 1.9% for the AstraZeneca–Oxford vaccines.

ARR (and NNV) are sensitive to background risk—the higher the risk, the higher the effectiveness—as exemplified by the analyses of the J&J’s vaccine on centrally confirmed cases compared with all cases:⁸ both the numerator and denominator change, RRR does not change (66–67%), but the one-third increase in attack rates in the unvaccinated group (from 1.8% to 2.4%) translates in a one-fourth decrease in NNV (from 84 to 64).

Lancet Microbe 2021
 Published Online
 April 20, 2021
[https://doi.org/10.1016/S2666-5247\(21\)00069-0](https://doi.org/10.1016/S2666-5247(21)00069-0)

See Online for appendix




Figure: RRR and NNV with 95% CI ranked by attack rate in the unvaccinated (placebo) group for five COVID-19 vaccines

The lower the NNV and the higher the RRR, the better the vaccine efficacy. Details are in the appendix (p 3). RRR=relative risk reduction. NNV=numbers needed to vaccinate. NIH=US National Institutes of Health.

There are many lessons to learn from the way studies are conducted and results are presented. With the use of only RRRs, and omitting ARR, reporting bias is introduced, which affects the interpretation of vaccine efficacy.¹⁰ When communicating about vaccine efficacy, especially for public health decisions such as choosing the type of vaccines to purchase and deploy, having a full picture of what the data actually show is important, and ensuring comparisons are based on the combined evidence that puts vaccine trial results in context and not just looking at one summary measure, is also important. Such decisions should be properly informed by detailed understanding of study results, requiring access to full datasets and independent scrutiny and analyses.

Unfortunately, comparing vaccines on the basis of currently available trial (interim) data is made even more difficult by disparate study protocols, including primary endpoints (such as what is considered a COVID-19 case, and when is this assessed), types of placebo, study populations, background risks of COVID-19 during the study, duration of exposure, and different definitions of populations for analyses both within and between studies, as well as definitions of endpoints and statistical methods for efficacy. Importantly, we are left with the unanswered question as to whether a vaccine with a given efficacy in the study population will have the same efficacy in another population with different levels of background risk of COVID-19. This is not a trivial question because transmission intensity varies between countries, affected by factors such as public health interventions and virus variants. The only reported indication of vaccine effectiveness is the Israeli mass vaccination campaign using the Pfizer–BioNTech product. Although the design and methodology are radically different from the randomised trial,² Dagan and colleagues¹¹ report an RRR of 94%, which is essentially the same as the RRR of the phase 3 trial (95%) but with an ARR of 0.46%, which translates into an NNV of 217 (when the ARR was 0.84% and the NNV was 119 in the phase 3 trial). This means in a real-life setting, 1.8 times more subjects might need to be vaccinated to prevent one more case of COVID-19 than predicted in the corresponding clinical trial.

Uncoordinated phase 3 trials do not satisfy public health requirements; platform trials designed to address

public health relevant questions with a common protocol will allow decisions to be made, informed by common criteria and uniform assessment. These considerations on efficacy and effectiveness are based on studies measuring prevention of mild to moderate COVID-19 infection; they were not designed to conclude on prevention of hospitalisation, severe disease, or death, or on prevention of infection and transmission potential. Assessing the suitability of vaccines must consider all indicators, and involve safety, deployability, availability, and costs.

We declare no competing interests.

Copyright © 2021 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY-NC-ND 4.0 license.

*Piero Olliaro, Els Torreele, Michel Vaillant
 piero.olliaro@ndm.ox.ac.uk

Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, Oxford OX3 7FZ, UK (PO); Institute for Innovation and Public Purpose, University College London, London, UK (ET); Competence Center for Methodology and Statistics, Luxembourg Institute of Health, Strassen, Luxembourg (MV)

- 1 Zimmer C, Corum J, Wee S-L. Covid-19 Vaccine Tracker. <https://www.nytimes.com/interactive/2020/science/coronavirus-vaccine-tracker.html> (accessed March 10, 2021).
- 2 Polack FP, Thomas SJ, Kitchin N, et al. Safety and efficacy of the BNT162b2 mRNA COVID-19 Vaccine. *N Engl J Med* 2020; **383**: 2603–15.
- 3 Baden LR, El Sahly HM, Essink B, et al. Efficacy and safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N Engl J Med* 2021; **384**: 403–16.
- 4 Voysey M, Clemens SAC, Madhi SA, et al. Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil, South Africa, and the UK. *Lancet* 2021; **397**: 99–111.
- 5 Logunov DY, Dolzhikova IV, Shcheblyakov DV, et al. Safety and efficacy of an rAd26 and rAd5 vector-based heterologous prime-boost COVID-19 vaccine: an interim analysis of a randomised controlled phase 3 trial in Russia. *Lancet* 2021; **397**: 671–81.
- 6 US Food and Drug Administration. Vaccines and Related Biological Products Advisory Committee meeting: FDA briefing document. Dec 10, 2020. <https://www.fda.gov/advisory-committees/advisory-committee-calendar/vaccines-and-related-biological-products-advisory-committee-december-10-2020-meeting-announcement> (accessed March 10, 2021).
- 7 US Food and Drug Administration. Vaccines and Related Biological Products Advisory Committee meeting: FDA briefing document. Dec 17, 2020. <https://www.fda.gov/advisory-committees/advisory-committee-calendar/vaccines-and-related-biological-products-advisory-committee-december-17-2020-meeting-announcement> (accessed March 10, 2021).
- 8 US Food and Drug Administration. Vaccines and Related Biological Products Advisory Committee meeting: FDA briefing document. Feb 26, 2021. <https://www.fda.gov/advisory-committees/advisory-committee-calendar/vaccines-and-related-biological-products-advisory-committee-february-26-2021-meeting-announcement> (accessed March 10, 2021).
- 9 Olliaro P. What does 95% COVID-19 vaccine efficacy really mean? *Lancet Infect Dis* 2021; published online Feb 17. [https://doi.org/10.1016/S1473-3099\(21\)00075-X](https://doi.org/10.1016/S1473-3099(21)00075-X).
- 10 Brown RB. Outcome reporting bias in COVID-19 mRNA vaccine clinical trials. *Medicina (Kaunas)* 2021; **57**: 199.
- 11 Dagan N, Barda N, Kepten E, et al. BNT162b2 mRNA COVID-19 vaccine in a nationwide mass vaccination setting. *N Engl J Med* 2021; published online Feb 24. <https://doi.org/10.1056/NEJMoa2101765>.

THE LANCET

Microbe

Supplementary appendix

This appendix formed part of the original submission. We post it as supplied by the authors.

Supplement to: Olliaro P, Torreale E, Vaillant M. COVID-19 vaccine efficacy and effectiveness—the elephant (not) in the room. *Lancet Microbe* 2021; published online April 20. [https://doi.org/10.1016/S2666-5247\(21\)00069-0](https://doi.org/10.1016/S2666-5247(21)00069-0).

Appendix

Table 1. Relative risk, relative risk reduction, absolute risk reduction and numbers needed to treat with 95% confidence intervals from publicly-available information.

Table 2. Calculations

Supplementary table 1.

Company	vaccine arms	Relative risk							Vaccine efficacy: relative risk reduction				Absolute risk reduction				Numbers needed to vaccinate			
		n	N	Risk	RR	SE(RR)	95%LL	95%UL	(RRR)	SE (RRR)	95%LL	95%UL	ARR	SE(ARR)	95%LL	95%UL	NNV	SE(NNV)	95%LL	95%UL
Pfizer/BioNTech	BNT162b2 mRNA	8	18198	0.4396	0.0497	0.7096	0.0245	0.1011	95.03%	0.14%	89.89%	97.55%	0.0084	0.0014	0.0070	0.0098	119	0.0007	102	143
	Placebo	162	18325	8.8404																
Moderna/NIH	mRNA-1273	11	14134	0.7783	0.0592	0.6078	0.0322	0.1087	94.08%	0.19%	89.13%	96.78%	0.0124	0.0019	0.0104	0.0143	81	0.0010	70	96
	Placebo	185	14073	13.1457																
Gamaleya	GamCovidVac	13	14094	0.9224	0.0903	0.6133	0.0489	0.1667	90.97%	0.29%	83.33%	95.11%	0.0093	0.0029	0.0063	0.0122	108	0.0015	82	158
	Placebo	47	4601	10.2152																
Johnson & Johnson	Ad26.COV2.S	116	19514	5.9445	0.3338	0.2092	0.2708	0.4115	66.62%	0.21%	58.85%	72.92%	0.0119	0.0021	0.0097	0.0140	84	0.0011	71	103
	Placebo	348	19544	17.8060																
AstraZeneca/Oxford	ChAdOx1 nCov-19	37	5807	6.3716	0.3316	0.3699	0.2291	0.4800	66.84%	0.41%	52.00%	77.09%	0.0128	0.0041	0.0088	0.0169	78	0.0021	59	114
	Placebo	112	5829	19.2143																

Data source:

Company	vaccine	source	ref.
Pfizer/BioNTech	BNT162b2 mRNA	table 2	2
Moderna/NIH	mRNA-1273	figure 4	3
Gamaleya	GamCovidVac	table 2	5
Johnson & Johnson	Ad26.COV2.S	table 10	8
AstraZeneca/Oxford	ChAdOx1 nCov-19	table 2	4

Supplementary Table 2.

	diseased	healthy	All
Exposed	a	b	n1
Not exposed	c	d	n2
All			

Relative risk : $RR = \frac{a/n_1}{c/n_2}$

Confidence intervals of RR : $95\%CI = e^{Ln(RR) \pm 1.96 * SE}$ where $SE = \sqrt{\frac{b/a}{n_1} + \frac{d/c}{n_2}}$

Relative Risk Reduction: $RRR = 1 - RR$

Confidence intervals of RRR : $95\%CI = 1 - 95\%CI RR$

Absolute Risk Reduction: $aRR = \frac{c}{n_2} - \frac{a}{n_1}$

Confidence Intervals of aRR: $95\%CI = aRR \pm 1.96 * SE$ where $SE = \sqrt{\frac{a/n_1 \times (1 - a/n_1)}{n_1} + \frac{c/n_2 \times (1 - c/n_2)}{n_2}}$

Number Needed to Vaccinate: $NNV = \frac{1}{aRR}$

Confidence Intervals of NNV: $95\%CI = \frac{1}{95\%CI aRR}$